

**A) Vertaisarvioidut tieteelliset julkaisut**

(7) Keva, O., Taipale, S., Hayden, B., Thomas, S. M., Vesterinen, J., Kankaala, P. & Kahilainen, K. 2021. Increasing temperature and productivity change biomass, trophic pyramids and community-level omega-3 fatty acid content in subarctic lake food webs. *Global Change Biology* 27(2): 282–296. DOI: 10.1111/gcb.15387

(6) Vesterinen, J., Keva, O., Kahilainen, K., Strandberg, U., Hiltunen, M., Kankaala, P. & Taipale, S. 2020. Nutritional quality of littoral macroinvertebrates and pelagic zooplankton in subarctic lakes. *Limnology and Oceanography* 9999: 1–17. DOI: 10.1002/lno.11563

(5) Strandberg, U., Vesterinen, J., Ilo, T., Akkanen, J., Melanen, M. & Kankaala, P. 2020. Fatty acid metabolism and modifications in *Chironomus riparius*. *Philosophical Transactions of the Royal Society B* 375 (1804), 20190643. DOI: 10.1098/rstb.2019.0643

(4) Vesterinen, J., Devlin, S.P., Syväranta, J., Jones, R.I. 2017. Influence of littoral periphyton on whole-lake metabolism relates to littoral vegetation in humic lakes. *Ecology* 98(12): 3074–3085. DOI:10.1002/ecy.2012

(3) Vesterinen, J., Devlin, S.P., Syväranta, J., Jones, R.I. 2017. Epiphytic bacteria make an important contribution to heterotrophic bacterial production in a humic boreal lake. *Aquatic Microbial Ecology* 79(3): 247–257. DOI: 10.3354/ame01835

(2) Vesterinen, J., Syväranta, J., Devlin, S.P., Jones, R.I. 2016. Periphyton support for littoral secondary production in a highly humic boreal lake. *Freshwater Science* 35(4):1235–1247. DOI: 10.1086/689032.

(1) Vesterinen, J., Devlin, S.P., Syväranta, J., Jones, R.I. 2016. Accounting for littoral primary production shifts a highly humic boreal lake towards net autotrophy. *Freshwater Biology* 61: 265–276. DOI: 10.1111/fwb.12700

**B) Opinnäytteet**

(3) Vesterinen, J. 2017. Littoral energy pathways in highly humic boreal lakes. Doctoral dissertation. Jyväskylä: University of Jyväskylä. Available at: <http://urn.fi/URN:ISBN:978-951-39-7038-3>

(2) Vesterinen, J. 2013. Littoral contribution to whole-lake primary production in highly humic Lake Mekkojärvi. Master's thesis. Jyväskylä: University of Jyväskylä. Available at: <http://urn.fi/URN:NBN:fi:jyu-201308172166>

(1) Vesterinen, J. 2010. Use and management plan for the Näsijärvi fishing area for the years 2010–2014. Bachelor's thesis. Turku: Turku University of Applied Sciences.